

PCT


WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61C 5/00, A61K 31/54, 33/14, 33/30, 33/40		A1	(11) International Publication Number: WO 98/17195
			(43) International Publication Date: 30 April 1998 (30.04.98)
(21) International Application Number: PCT/US97/19598 (22) International Filing Date: 23 October 1997 (23.10.97) (30) Priority Data: 08/736,356 23 October 1996 (23.10.96) US (71) Applicant (for all designated States except US): THE RESEARCH FOUNDATION OF STATE UNIVERSITY OF NEW YORK [US/US]; State University of New York at Stony Brook, Stony Brook, NY 11794-3366 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): KLEINBERG, Israel [US/US]; 14 Three Pond Road, Smithtown, NY 11787 (US). CODIPILLY, Milroy [LK/US]; 305A Chelsea Place, Coram, NY 11727 (US). (74) Agents: RYAN, Charles, S. et al.; Darby & Darby P.C., 805 Third Avenue, New York, NY 10022 (US).			(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report.

(54) Title: COMPOSITIONS TO CONTROL ORAL MICROBIAL OXIDATION-REDUCTION (E_h) LEVELS



(57) Abstract

The present invention relates to an oral composition containing a zinc compound containing free available zinc ion and at least one stabilized or stable E_h raising compound distributed in an oral vehicle. The present invention further relates to a method of inhibiting the formation of sulfur containing anions and preventing a reduction in the E_h of the oral cavity. A method of reducing oral malodor and gingivitis and periodontitis is also provided by this invention.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AX	Åland Islands	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BD	Bangladesh	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BE	Belgium	GR	Greece	ML	Mali	TR	Turkey
BF	Burkina Faso	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BS	Bahamas	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

5 COMPOSITIONS TO CONTROL ORAL MICROBIAL
OXIDATION-REDUCTION (E_h) LEVELS

FIELD OF INVENTION

10 The present invention provides oral compositions comprising a zinc compound containing free available zinc and at least one stabilized or stable E_h raising compound distributed in an oral vehicle. The present invention is further directed to a method of inhibiting the formation of anionic sulfur species in the oral cavity and preventing a lowering of the E_h of the oral cavity. A method of simultaneously reducing
15 oral malodor, gingivitis and periodontitis by preventing or reducing oral putrefaction is also provided by this invention.

BACKGROUND OF THE INVENTION

The hard and soft tissues of the mouth are covered with microbial
20 populations that contain bacteria with different metabolic capabilities. The Gram-positive bacteria within these microbial populations readily catabolize carbohydrates to produce acids which attack the hard tissues of the oral cavity, resulting in the formation of dental caries lesions (cavities). In contrast, the Gram-negative bacteria, especially the anaerobes readily metabolize various amino acids contained in salivary (and to lesser extent other)
25 peptides and proteins in the oral cavity to form end-products which favor the formation of oral malodor and periodontitis. This process of peptide, protein and amino acid degradation by the mouth bacteria is referred to as oral bacterial putrefaction. The mixture of malodorous compounds produced by the Gram-negative anaerobic bacteria during putrefactive degradation of proteins, peptides and amino acids include hydrogen sulfide,
30 methyl mercaptan, and dimethyl sulfide (formed from the sulfur containing amino acids cysteine, cystine and methionine); indole and skatole (formed during the metabolism of tryptophan); cadaverine and putrescine (produced from lysine and ornithine); and butyrate and valerate (produced from the metabolism of other amino acids). The production of these malodorous compounds in the oral cavity results in a condition commonly referred to as
35 oral malodor.

Hydrogen sulfide, methyl mercaptan, butyrate and propionate are putrefaction end-products that also have cell and tissue altering non-inflammatory roles in the periodontitis process. Hydrogen sulfide and methylmercaptan are compounds particularly effective in facilitating the oral epithelium penetrability of toxins and other large molecular weight compounds produced by Gram-negative bacteria, and leading to the inflammation and tissue degradation characteristics of gingivitis and periodontitis. Gingivitis is a condition in which the gingiva is red, swollen and bleeding. If left untreated, gingivitis may develop into periodontitis, a condition characterized by destruction of the periodontium, including epithelial attachment loss, periodontal membrane and ligament destruction, and loss of gingiva and alveolar bone. Severe periodontitis resulting in deep periodontal pockets may ultimately result in tooth loss.

Previous studies have largely focused on the use of germicidal agents to treat gingivitis-periodontitis and oral malodor. Until the findings by the present invention, previous studies have not recognized that gingivitis-periodontitis and oral malodor arise from a common process, oral bacterial putrefaction; also that this putrefaction can be inhibited by simultaneously lowering the ability of the oral bacteria to reduce the oxidation-reduction potential (E_h) of the oral cavity and at the same time, raising the existing E_h to where the oral environmental E_h is not conducive to oral putrefaction and oral disease production.

The metabolism and growth of anaerobic bacteria in the oral cavity is favored when the E_h is lowered. The present invention has discovered that the oxidation-reduction potential (E_h) is a key regulating factor in oral bacterial putrefaction. The lowering of the E_h of the oral cavity has been found to occur in two steps, a depletion of oxygen followed by the generation of electron rich compounds. The present invention has discovered from studies on the isolation and catabolism by the oral bacteria of nitrogenous substrates in human saliva that the primary compounds responsible for lowering the E_h of the oral cavity are non-volatile sulfur containing anions derived largely from cysteine and cystine (Table 1). These include the anionic sulfur species, sulfide (S^{2-}), hydrogen sulfide (HS^-) and methyl mercaptan (CH_3S^-). Such anions favor an ecological environment of reduction (lower E_h) that enables the Gram-negative anaerobic bacteria in the mouth to grow, engage in oral putrefaction and produce electron-rich compounds leading to and maintaining a prolonged lowering of the E_h of the oral cavity and the undesirable

conditions of oral malodor, gingivitis and periodontitis. A lower E_h favors the oral bacterial putrefaction process whereas a higher E_h is inhibitory.

Table 1. Lowest E_h reached with the common amino acids when incubated with the mixed bacteria in salivary sediment.

amino acids	E_h (millivolts)		
	Group A	Group B	Group C
alanine	100		
arginine		25	
asparagine	70		
aspartic	80		
cysteine			20
cystine			-10
glutamic		30	
glutamine	100		
histidine	70		
isoleucine	115		
leucine	110		
lysine	95		
methionine			20
ornithine		55	
phenylalanine	100		
proline	100		
serine	115		
threonine	110		
tryptophan	85		
tyrosine		40	
valine	105		
salivary supernatant	-25; water control	142	

The present invention provides compounds that simultaneously (i) inhibit the formation of these electron-rich compounds and thereby prevent the E_h from falling to harmful levels and (ii) react with any electron-rich compounds formed and by thus neutralizing them, raise the E_h to safer levels. It has been surprisingly discovered in accordance with the present invention that an oral composition containing a zinc compound capable of providing free available zinc and a stabilized or stable E_h raising compound can effectively prevent the lowering of the E_h . This is crucial to preventing oral bacterial putrefaction, the metabolic

process that is the basis and that results in the development of both oral malodor and gingivitis-periodontitis.

Zinc compounds, hydrogen peroxide, and chlorine dioxide have each been used as therapeutic agents in oral compositions to destroy harmful bacteria involved in oral malodor and gingivitis-periodontitis formation. Previous studies where zinc has been identified as having anti-microbial and anti-plaque effects have made no distinction between zinc compounds where the zinc is freely available and where it is not. (See for example, U.S. Patent No. 4,289,755 to Dhabhar). The zinc species is an important element of this invention, since free available zinc (zinc ion) within the oral cavity is required to inhibit the E_h lowering capability of a pathogenic, putrefactive microbiota. Zinc compounds used in the compositions previously described include compounds where zinc is not freely available. Zinc that is bound or complexed to various ligands and zinc species that have a low solubility and form precipitates are prevented from reacting with the E_h lowering enzymes and products produced by the putrefactive microbiota and hence are poorly suited for the purposes of this invention.

The solubility for zinc compounds vary as shown in the table below.

Solubility of selected zinc compounds		
Compound	Formula	Solubility (g/100 cc)
zinc chloride	$ZnCl_2$	432 @ 25°C
zinc citrate	$Zn_3(C_6H_5O_7)_2$	slightly soluble
zinc acetate	$Zn(C_2H_3O_3)_2$	30 @ 20°C
zinc lactate	$Zn(C_3H_5O_3)_2$	5.7 @ 15°C
zinc salicylate	$Zn(C_7H_5O_3)_2$	5 @ 20°C
zinc sulfate	$ZnSO_4$	soluble
zinc oxide	ZnO	0.00016 @ 29°C
zinc nitrate	$ZnNO_3)_2$	infinitely soluble

Data are from the *Handbook of Chemistry and Physics*, Chemical Rubber Company, 67th Edition CRC Press, Boca Raton, Florida, 1986-87.

It is evident from this table that the amount of zinc that will be soluble and available in the oral cavity for controlling pathogenic microbiota will vary considerably. Those zinc compounds that provide low levels of zinc ion in solution, such as zinc oxide, are

unsuitable for the present invention. This distinction between the various zinc compounds was not recognized prior to this invention. In essence, in order to inhibit the E_h from falling to harmful levels, it is essential that the zinc ions are freely available.

Previous studies have also identified hydrogen peroxide and chlorine dioxide as
5 germicidal agents. The chlorine dioxide is usually derived from the chlorite ion.
Hydrogen peroxide owes its germicidal activity to oxygen release and formation of free radicals which provides chemical and mechanical mechanisms for killing oral anaerobic bacteria and cleaning wounds and removing tissue and other debris from inaccessible areas (such as between the teeth). The release of oxygen from hydrogen peroxide is particularly pronounced in the
10 presence of catalase, organic matter, metals and metal compositions. In this invention, this is inhibited from happening by use of chloride ion, so that the peroxide will serve as an E_h raising compound rather than as a germicidal agent. U.S. Patent Nos. 5,104,644, 5,174,990 and 5,310,546 to Douglas describe the use of hydrogen peroxide as such a germicidal agent. Specifically in the oral composition described by Douglas, hydrogen peroxide releases
15 molecular oxygen in the presence of tissue catalase and peroxidase to act against the oral anaerobic bacteria. Previous studies do not stabilize the hydrogen peroxide so that it does not break down.

The oral composition described herein acts as an E_h raising compound rather than one that results in formation of breakdown products. In this invention, degrading these
20 species to produce a germicidal effect, is avoided. The hydrogen peroxide in the composition described herein is stabilized using chloride ion, an acidic pH, and avoiding mixing it with zinc ion until just before use.


Chlorine dioxide is an oxyhalogen compound widely used in industry for disinfection and control of bacterial biofouling. It is also used to control taste, odor, oxidation
25 of metal ions and color removal in other applications. Several studies have described using chlorine dioxide as an antimicrobial agent in mouth rinse applications. For example, U.S. Patent No. 4,696,811 to Ratcliff describes a method and composition to destroy malodorous compounds; Patent No. UK 2290233A to Drayson and Butcher describes compositions for tooth whitening. Others include inventions where the oxidizing and germicidal capabilities of chlorine
30 dioxide are activated by forming chlorine dioxide just prior to use. The main reason chlorine dioxide is generated in this way is because chlorine dioxide is an unstable gas at room temperature (boiling point of 11°C) and is sensitive to decomposition by visible and ultraviolet light. In previous studies, the chlorine dioxide is commonly generated from the chlorite ion by

acidification. It is usually provided as sodium chlorite buffered to a pH around 7 to 8 and above, and as such, referred to as stabilized chlorine dioxide (U.S. Patent Nos. 4,689,215 and 4,837,009 to Ratcliff, Patent No. UK 2290233A to Drayson and Butcher and Patent No. WO 95/27472 to Richter). Chlorine dioxide like hydrogen peroxide above, is typically generated for the purpose of killing mouth bacteria. Where acidification of chlorite is carried out, the levels of chlorine dioxide produced are usually adequate for its germicidal purpose. But in those compositions where chlorine dioxide is stabilized as sodium chlorite at neutral or alkaline pH, formation of chlorine dioxide from chlorite ion is a relatively slow process. Accordingly, very little chlorine dioxide is available within the oral cavity as an antibacterial agent in these compositions. In contrast, in the compositions herein, chlorite ions are not used as germicidal agents. Instead they are used as effective and stable E_h raising compounds, if their degradation to chlorine dioxide is avoided.

In previous inventions, chlorite ion use involves provision (rather than removal) of electrons to enable chlorite ion disproportionation and formation of the bacteriocidal compound, chlorine dioxide, a process stimulated by acid addition. The oxidation-reduction change involves change in oxidation state from +3 to +4. On the other hand, when chlorite acts as an E_h raising compound as in the invention herein, its oxidation state decreases from +3 to -1. Reduction of chlorite ion is to chloride ion after going through a series of reactions involving various intermediates. What is evident from this analysis is that the chlorite ion is capable of acting as either an oxidizing or as a reducing agent depending upon the reaction conditions. Few compounds show such multi-step, and hence atypical redox buffer effects. Nonetheless, this enables them to counter or resist along with zinc ion the kinds of changes in the E_h level that enable oral putrefaction to flourish and be suitable for this invention.

Using sodium chlorite as an E_h raising compound rather than as a source of chlorine dioxide is very important, because chlorine dioxide at elevated levels combines with certain amino acids to produce compounds that are potentially mutagenic. Therefore, inhibition or prevention of significant chlorine dioxide formation from sodium chlorite is desirable and preferred and contra-indicated is the utilizing of sodium chlorite to generate large amounts of chlorine dioxide therefrom in order to kill enough bacteria to have significant oral effects.

A neutral pH and above is essential for chlorite ion stability and to avoid chlorine dioxide formation. Also, chloride ion is useful for additional stabilization of the sodium chlorite where there is any decrease in the pH. This is because chloride ion is produced when chlorite becomes chlorous acid and disproportionation of chlorous acid occurs



This reaction is inhibited by mass action when chloride ion is provided.

Hydrogen peroxide like sodium chlorite behaves as an E_h raising compound in that it can also readily react as either an oxidizing or a reducing agent. Analogous to the chlorine dioxide/chlorite/chloride system is the molecular oxygen/hydrogen peroxide/water system where chlorite ion and hydrogen peroxide are similarly positioned as oxidation-reduction intermediates. In previous studies, the peroxide is used to generate oxygen and/or oxygen reactive species to kill the oral bacteria involved in oral disease. As found for chlorite ion, peroxide in this invention functions as an E_h raising compound by removing the excess electrons of putrefaction and producing hydroxyl ions rather than giving up electrons and undergoing disproportionation as in previous patents.

The disproportionation reaction of hydrogen peroxide



is very slow under ordinary conditions but rapid in the presence of the enzymes, catalase and peroxidase, found in certain of the bacteria in the oral cavity. Once again, hydrogen peroxide is a species like chlorite ion that is thermodynamically unstable with respect to disproportionation and can function as an intermediate in oxidation-reduction reactions. Further and once again, the chloride ion inhibits the disproportionation of hydrogen peroxide but does so through inhibition of catalase.

The oxygen-oxygen single bond in hydrogen peroxide is one of the weakest covalent bonds known. It is easily broken indicating that it readily accepts electrons and as a result is able to produce hydroxyl ions. Alternatively, hydrogen peroxide is converted into the stable oxygen molecule. Previous studies have shown, using ^{18}O labeled hydrogen peroxide, that the oxygen produced is derived entirely from the peroxide species and not from water. This suggests that the breakdown of peroxide does not involve the breaking of the O-O bond but provides electrons to an appropriate oxidizing agent. H_2O_2 when used as an E_h raising compound is not used as a source for molecular oxygen.

Recent studies by Douglas (Patents Nos. 5,104,644, 5,174,990, and 5,310,546) have described oral compositions combining zinc chloride and hydrogen peroxide to treat gingivitis-periodontitis. U.S. Patent No. 5,174,990 describes a mouth rinse containing zinc

chloride and hydrogen peroxide. In these Douglas patents, it is necessary to counter the instability prior to use of the formulations described, which is caused largely because of the presence of zinc metal. To aid in stabilization of zinc, ligands that bind well to zinc such as citrate and laurylsulfate are added. However, these additions reduce the availability of free zinc, especially when these ligands are present at high ligand to zinc ratios. The mouth rinses in Patent Nos. 5,174,990 and 5,310,546 have zinc chloride concentration ranging from 0.005% to 0.1% and a hydrogen peroxide concentration ranging from 0.25% to 0.65%. In the absence of ligands that bind zinc, zinc ion at these zinc chloride levels range in concentration between 0.002% and 0.047%. Moreover, recent studies have shown that hydrogen peroxide at the concentrations described therein are degraded by oral bacterial catalase and are not effective *in vivo*. Ryan and Kleinberg (1995) *Archs. oral Biol.*, 40, 753-763. Accordingly, to compensate for the rapid breakdown of hydrogen peroxide by catalase, use of higher concentrations of hydrogen peroxide (1% or above) is necessary. At hydrogen peroxide concentrations above 3.0 to 3.5%, studies have shown that hydrogen peroxide can be harmful to the soft tissues of the oral cavity. Thus, for an oral composition containing hydrogen peroxide to be effective as a therapeutic and at the same time not harm the soft or hard tissues of the oral cavity or be harmful if swallowed, its concentration needs to be between about 1% and 3%.

In contrast, the oral compositions discovered in this invention contain sufficient chloride ions to inhibit catalase hydrolysis of the peroxide, thereby enabling peroxide to remain intact even at lower peroxide levels than 1% and for the peroxide to serve as an E_h raising compound where formation of toxic products from peroxide are avoided.

Many reactions involving hydrogen peroxide in solution result in the production of free radical species, such as HO_2 and OH . These are effective agents for killing bacteria and such formation is a basis for the use of hydrogen peroxide as a disinfectant. Transition metal-ion catalyzed decomposition of hydrogen peroxide can readily give rise to the formation of free radicals. The destabilizing effects of zinc are avoided in this invention by using a two compartment approach where combination with zinc compound is provided just prior to use to ensure maximum availability of free zinc. The compositions described herein result in a synergism between the zinc ions, the peroxide used as an E_h raising compound and the chloride ions. This two compartment system is a more desirable and effective approach than oral compositions previously described.

Methylene blue has been used as an E_h raising compound. It readily accepts electrons and in this way helps prevent the electron accumulation that produces the low E_h that

favors oral putrefaction. In U.S. Patent No. 5,087,451 to Wilson and Harvey, methylene blue is used to inhibit periodontitis. The beneficial effect of methylene blue alone is significantly less than achieved when used in conjunction with zinc ion as with the compositions of this invention.

5

SUMMARY OF THE INVENTION

The present invention is directed to oral compositions containing a zinc compound where zinc ion is freely available and at least one stabilized or stable E_h raising compound distributed in an oral vehicle. In one embodiment, the oral compositions contain a
10 zinc compound containing freely available zinc ions, hydrogen peroxide and a chloride ion containing compound. In another embodiment, the oral composition contains a zinc ion compound and an oxyhalogen compound. A further embodiment of this invention includes an oral composition containing a zinc ion compound and methylene blue.

The present invention is further directed to a method of inhibiting the formation
15 of anionic sulfur species and thereby preventing a lowering of the E_h of the oral cavity. Specifically, the method comprises delivering into the oral cavity a therapeutically effective amount of an oral composition containing a free available zinc compound and at least one stabilized or stable E_h raising compound distributed in an oral vehicle.

A further embodiment of the present invention is a method of reducing both oral
20 malodor and gingivitis-periodontitis comprising delivering into the oral cavity a therapeutically effective amount of an oral composition containing a zinc ion compound and at least one stabilized or stable E_h raising compound.

BRIEF DESCRIPTION OF THE DRAWINGS

25 Figure 1 shows the relation between zinc ion concentration and the rate at which the oral bacteria are able to produce volatile sulfur compounds and associated sulfur anions from cysteine. Inhibition is exponential and occurs maximally once the zinc ion concentration reaches about 0.2%.

Figure 2 shows the changes in the E_h during an incubation of mixed oral bacteria
30 and (i) salivary supernatant (ii) glucose (iii) salivary supernatant and glucose and (iv) a water control.

Figure 3 shows the changes in E_h during an incubation of mixed oral bacteria and salivary supernatant with (i) zinc chloride (ii) hydrogen peroxide and (iii) zinc chloride ion

plus hydrogen peroxide. A control incubation containing mixed oral bacteria with salivary supernatant is also shown. Hydrogen peroxide by itself loses some of its ability to inhibit the E_h lowering due to hydrogen peroxide degradation by bacterial catalase during the incubation.

Figure 4 shows the mean E_h , odor index, volatile sulfur compound (VSC) level and the level of indole/skatole produced in samples taken at intervals in a 4 hour period after rinsing with a mouth rinse containing zinc chloride (0.08%), hydrogen peroxide (1%) and NaCl (2.9%). VSC levels are measured with an instrument called the Halimeter.

Figure 5 shows the concentration of volatile sulfur compounds and the E_h *in vivo* following rinsing with cysteine (VSC and E_h responses) before and after a mouth rinse containing zinc chloride at 6 mM (0.08%).

Figure 6 shows the concentration of volatile sulfur compounds and the E_h *in vivo* following rinsing with cysteine (VSC and E_h responses) before and after a mouth rinse containing zinc chloride at 6 mM (0.08%), hydrogen peroxide at 1% and sodium chloride at 500 mM (2.9%).

Figure 7 shows the concentration of volatile sulfur compounds and the E_h *in vivo* following rinsing with cysteine (VSC and E_h responses) before and after a mouth rinse containing zinc chloride at 6 mM (0.08%) and sodium chlorite at 0.5%.

Figure 8 shows the concentration of volatile sulfur compounds *in vivo* following rinsing with cysteine (VSC and E_h responses) before and after a mouth rinse containing sodium chlorite at 0.1%.

Figure 9 shows the concentration of volatile sulfur compounds *in vivo* following rinsing with cysteine (VSC and E_h responses) before and after a mouth rinse containing sodium chlorite at 0.1% and zinc chloride at 6 mM (0.08%).

DETAILED DESCRIPTION OF THE INVENTION

The essential components and their relevant proportions in the compositions of the invention are set forth below. All patents, publications and test methods mentioned herein are incorporated by reference.

The present invention relates to an oral composition containing a zinc ion compound where a high concentration of free available zinc is provided and at least one E_h raising compound distributed in an oral vehicle. A zinc ion compound as defined by the present invention is a compound containing freely available zinc ions capable of inhibiting the lowering of the oral cavity E_h . Important in this regard is inhibiting the breakdown of cysteine

or cystine from saliva, mucosal tissues (especially the tongue) and foods by the oral bacteria. Freely available zinc ions are ions that are not bound. Zinc compounds present in the oral compositions of this invention include, for example, zinc chloride, zinc acetate, zinc lactate, zinc salicylate, zinc sulfate, and zinc nitrate. In one embodiment of the present invention, the zinc
5 compound is zinc chloride. The concentration of zinc ion in the oral composition may range from about 0.02% to about 0.2%. In a preferred embodiment, the zinc ion concentration ranges from about 0.04% to about 0.12%.

A stabilized E_h raising compound as defined herein is a compound capable of directly or indirectly raising the E_h of the oral cavity. Examples of E_h raising compounds
10 include, for example, hydrogen peroxide, oxyhalogen species such as sodium chlorite and biologically compatible oxidation-reduction (redox) buffers such as methylene blue. Additional examples of E_h raising compounds include common fermentable sugars such as glucose, galactose, fructose, maltose, lactose and sucrose. These compounds when metabolized by oral bacteria and in particular, the oral streptococci, in the presence of oxygen produce, *inter alia*,
15 hydrogen peroxide.

The concentration of the E_h raising compound in the oral compounds of this invention may range from about 0.1% to about 3.0% by weight of the composition. In a preferred embodiment, the concentration of the E_h raising compound is about 0.1% to about 1.0%. When the E_h raising compound is hydrogen peroxide or a fermentable sugar, a chloride
20 ion containing compound is added to the oral composition in an amount sufficient to inhibit catalase in the oral cavity from breaking down the hydrogen peroxide. The chloride ion containing compound is a compound capable of inhibiting the catalase activity in the oral cavity. Suitable chloride ion containing compounds include alkali metal chloride salts and alkali earth metal chloride salts such as, for example, NaCl and $CaCl_2$. Generally, the
25 concentration of the chloride ion containing compound in the oral composition ranges from about 0.5% to about 2.5% by weight of the composition. Since some oral microbiota produce hydrogen peroxide in the presence of fermentable sugars, chloride ions are also added to make it more effective.

In one embodiment of this invention the zinc compound is zinc chloride, the E_h
30 raising compound is hydrogen peroxide and the chloride ion containing compound is sodium chloride. In another embodiment of this invention, the oral composition contains zinc chloride, a fermentable sugar and sodium chloride. In a further embodiment of this invention, the oral composition contains zinc chloride and the oxyhalogen species sodium chlorite and with or

without sodium chloride. A further embodiment of the present invention consists of oral compositions containing zinc chloride and methylene blue with or without sodium chloride.

We have discovered that the pH of the oral compositions of the present invention is specific to the zinc compound and E_h raising compound combination used. When the E_h raising compound is hydrogen peroxide, the preferred pH of the oral compositions generally ranges from about 3.0 to about 6.0. In one embodiment, the pH ranges from about 3.5 to about 4.5. An acidic pH has two desirable effects. First, an acidic pH ensures availability of the zinc ion since zinc ion above a pH of about 6.0 combines with hydroxyl ions in solution to form poorly soluble zinc hydroxide, thereby making zinc ion unavailable. Second an acidic pH converts sulfur anions to the acidic forms which result in a higher E_h . In the case of hydrogen sulfide, because hydrogen sulfide is volatile, its formation serves as an effective means of getting rid of electrons carried by sulfide anion that are particularly conducive to lowering the E_h . Third, catalase degradation of hydrogen peroxide is inhibited at an acidic pH. An acidic pH and the presence of chloride ion, ensures that the hydrogen peroxide in the composition is not degraded with storage and hence retains its effectiveness.

On the other hand, when the E_h raising compound is the oxychloride, sodium chlorite, a pH between about 3.0 and 6.0 is unsuitable for the stability of sodium chlorite during its storage. At acidic pH, unstable and less desirable chlorine dioxide is produced. To be of use for the purpose of this invention, the pH of the sodium chlorite during storage needs to be between about 7.0 and about 8.5 where it is most stable. The instability of zinc ion at a pH of 6.0 and above and the instability of chlorite at a pH of about 6.0 and below necessitates the two being kept separate in a two compartment system until ready for use. When brought into contact with each other immediately before mixture use, the preferred pH is between about 5.5 and about 6.0. Addition of chloride ion as sodium chloride and/or as part of zinc chloride, provides stability along with synergistic activity. The pH of the oral compositions described herein can be controlled with acids such as hydrochloric and benzoic and with base such as sodium hydroxide.

In addition to the zinc-peroxide-chloride ion composition, the zinc-oxyhalogen composition, and the zinc-methylene blue composition, the oral compositions described in accordance with the present invention may contain any conventional ingredient for the particular oral composition. For example, liquid mouthwashes may contain a solvent such as distilled or deionized water, and ethanol; a sweetening agent such as sorbitol, mannitol, xylitol, saccharin and aspartame; and a flavoring agent such as peppermint oil, and spearmint oil. (See U.S.

Patent Nos. 4,226,851; 4,209,754; 4,289,755; and 5,104,644). Because of the formulation difficulties associated with zinc ion instability at about pH 6.0 and above resulting from the forming of poorly soluble zinc hydroxide and the instabilities of many E_h raising compounds such as sodium chlorite and hydrogen peroxide, two compartment systems where the zinc ion compound and the E_h raising compound are in respective compartments but can be mixed immediately prior to use are preferred. An example of a two compartment composition for zinc chloride and sodium chlorite is shown in Table 2.

Dentifrices may contain, for example, a conventional abrasive such as calcium pyrophosphate, aluminum hydroxide, resins, insoluble alkali metal metaphosphates and silica in a standard amount of 20-60% wt.; a binder such as hydroxyethyl cellulose, xanthin gum and sodium carboxymethylcellulose in a standard amount of 0.5-5.0% wt.; a foaming agent such as sodium lauryl sulfate, sodium coconut monoglyceride sulfonate and sodium-N-methyl-N-palmitoyl tauride in a standard amount of 0.5-3.0% wt.; a flavoring agent; a sweetening agent; an antiseptic agent and any other ingredient required for the particular formulation. (See U.S. Patent No. 5,372,802). Two compartment delivery systems are preferred. Tablets and powders may contain, for example, a vehicle such as lactose or mannitol; a binder such as corn starch or carboxymethyl cellulose; and a disintegrator, once more in two compartment delivery systems.

The present invention is also directed to a method of inhibiting the formation of sulfur containing anions in the oral cavity and preventing a lowering of the E_h of the oral cavity by delivering into the oral cavity a therapeutically effective amount of an oral composition containing, a zinc ion compound and at least one E_h raising compound distributed in an oral vehicle. As defined by the present invention, sulfur containing anions include for example, sulfide (S^{2-}), hydrogen sulfide anion (HS^-) and methyl mercaptan anion (CH_3S^-). A therapeutically effective amount of oral composition is an amount sufficient to inhibit the formation of sulfur containing anions and prevent a lowering of the E_h of the oral cavity. For example, a therapeutically effective amount of the oral composition in a dentifrice or mouth rinse may range from approximately 0.5% to approximately 5% by weight and preferably 2% to 3% by weight of the composition.

Acceptable oral vehicles include, for example, any conventional oral delivery system, such as dental care products, food products and chewing gum. Examples of dental care products include for example, dentifrices, topical solutions or pastes, mouthwashes in the form of liquids, powders, gels or tablets, and dental flosses. Examples of food products which

contain the oral compositions described herein include, for example, lozenges and confectioneries.

The present invention is further directed to a method of reducing oral malodor, gingivitis and periodontitis by delivering into the oral cavity a therapeutically effective amount of an oral composition containing a zinc ion compound and at least one E_h raising compound distributed in an acceptable oral vehicle. As defined by the present invention, a therapeutically effective amount of an oral composition is an amount sufficient to raise the E_h of the oral cavity to normal levels and prevent or reduce oral malodor, gingivitis and periodontitis. For example, a therapeutically effective amount of an oral composition is an amount sufficient to reduce or prevent the formation of malodorous compounds such as hydrogen sulfide and the growth of harmful Gram-negative anaerobic bacteria which may cause gingivitis and periodontitis. A therapeutically effective amount of the oral composition in a dentifrice or mouthwash may range from approximately 0.5% to approximately 5% by weight and preferably about 2% to about 3% by weight of the composition.

The present invention further provides an article of manufacture comprising a packaging material and the oral compositions described herein contained within said packaging material, wherein said oral composition is effective in preventing and/or reducing any decrease in the E_h , oral putrefaction and development of oral malodor, gingivitis and periodontitis and wherein said packaging material contains a label that indicates that said oral composition is effective in raising the E_h , and reducing oral putrefaction, oral malodor, gingivitis and periodontitis. The packaging material used to contain the oral compositions can comprise glass, plastic, metal or any other suitably inert material. For example, a dentifrice containing the oral composition of the present invention may be contained in a collapsible tube, typically aluminum, lined lead or plastic or a squeeze, pump or pressurized dispenser for measuring out the contents or in a tearable sachet.

In order to further illustrate the present invention, the experiments described in the following examples were carried out. It should be understood that the invention is not limited to the specific examples or the details described therein. The results obtained from the experiments described in the examples are shown in the accompanying table and figures.

Example I

This Example demonstrates the ability of salivary supernatant to lower the E_h and the ability of glucose to act as an E_h raising compound to raise the E_h in the salivary

sediment system model developed by Kleinberg (1967 and 1970) *Archs. oral Biol.* 12:1457-1473; *Advances oral Biol.* (Volume 4) New York, NY Academic Press pp. 49-50 (see Figure 2). This model utilizes the mixed oral bacteria collected in the salivary sediment obtained from whole saliva by centrifugation. Extensive studies have shown that this system behaves metabolically like dental plaque (Singer *et al.* (1983) *Archs. oral Biol.* 28:29-35; Wijeyeweera and Kleinberg (1989 a and b) *Archs. oral Biol.* 34:43-64; Salako and Kleinberg (1992) *Archs. oral Biol.* 37:821-829; Ryan and Kleinberg (1995) *Archs. oral Biol.* 40:743-752). This model is recognized in the art as an effective proxy to study the microbial metabolic and related biochemical processes of the oral cavity. The ability of the different amino acids available from salivary peptides and proteins to lower the E_h in the salivary sediment system is shown in Table 1.

To collect salivary sediment and salivary supernatant for the assay, saliva stimulated by chewing paraffin wax was expectorated into test-tubes chilled in cracked ice. The donors had not brushed their teeth for 24 hours prior to collection. The donors were also instructed to fast for at least 12 hours to provide stimulated whole saliva which contains a minimal level of endogenous carbohydrate. (Kleinberg and Jenkins (1964) *Archs. oral Biol.* 9:493-516). Following collection, the saliva was centrifuged at 1740 X g for 15 minutes and the supernatant was subsequently removed by pipette and kept on ice until the assay was run. Next, the sediment was washed three times with ice cold distilled water to remove any residual salivary supernatant. The washed salivary sediment was chilled in crushed ice until the assay was run. Prior to incubating the samples and running the assay, the washed salivary sediment was re-suspended in distilled water to a final concentration of 50 per cent (V/V).

To assay the ability of various amino acids and salivary supernatant to produce a lowering of the E_h , incubation tubes were prepared containing the following: 16.7% (V/V) salivary sediment, 60 mM phosphate buffer and either 33.3% (V/V) salivary supernatant or 3 mM amino acid. A control tube contained sediment and water. All incubation mixture preparations were done with the test-tubes chilled in cracked ice until they were transferred to a water bath for incubation at 37°C for 24 hours. Measurements were taken at the following time intervals: 0 and 30 min and 1, 2, 3, 8 and 24 hours.

The oxidation-reduction potential (E_h) was determined in each incubation mixture using a platinum electrode and a potassium chloride salt bridge leading to a calomel reference electrode connected to a pH meter used as a millivoltmeter. All E_h measurements were related to the standard hydrogen electrode by adding the value 242 millivolts to the

reading made with this electrometric system. The lowest E_h reached with each amino acid is shown in Table 1. The corresponding lowest E_h levels reached with salivary supernatant and the water control are shown at the bottom of the table. The E_h with supernatant falls to the lowest E_h and is only matched by the amino acid, cystine. Of the various amino acids tested, the sulfur containing amino acids - cystine, cysteine and methionine (Group C) showed the lowest E_h ; arginine, glutamic, ornithine and tyrosine (Group B) showed next lowest; and the remainder (Group A) showed the highest E_h levels. Fractionation experiments of salivary supernatant identified small peptides with cystine as the constituent mainly responsible for the E_h lowering activity of salivary supernatant.

Example II

A similar type of *in vitro* experiment as in Example I was carried out to assay the ability of (i) zinc chloride, (ii) hydrogen peroxide, and (iii) zinc chloride and hydrogen peroxide, to prevent a fall in the E_h of the oral cavity. Incubation tubes were prepared containing 16.7% sediment, 60 mM phosphate buffer, and (i) zinc chloride at a concentration of 6.0 mM (0.08%), (ii) hydrogen peroxide at a concentration of 0.5%, or (iii) zinc chloride at a concentration of 6.0 mM and hydrogen peroxide at a concentration of 0.5%. A tube was prepared containing salivary sediment and salivary supernatant only (water control). Figure 2 shows the E_h of the mixtures during incubation at 37°C over a 24-hour time period.

As shown in Figure 3, zinc chloride and hydrogen peroxide alone, or in combination, significantly raised the E_h of the incubation mixture compared to the salivary supernatant plus salivary sediment control. The control incubation mixture containing only sediment and supernatant showed a rapid and extensive lowering of the E_h . The incubation mixtures with zinc chloride alone and hydrogen peroxide alone, showed some decrease in the E_h . Hydrogen peroxide alone progressively lost some of its E_h reducing ability as there was no chloride present unlike in zinc chloride to inhibit its degradation by catalase. When zinc chloride and hydrogen peroxide were combined in the incubation, only a slight decrease in the E_h occurred. The results in this example demonstrated the ability of a composition containing zinc chloride and hydrogen peroxide to reduce the decrease in the E_h , and thereby reduce the oral putrefaction that is conducive to the development of oral malodor, gingivitis and periodontitis.

Example III

This example shows the ability of the zinc ion-peroxide-chloride ion compositions of the present invention to retard or reduce the E_h lowering ability of the oral bacteria and their ability to produce malodor. As shown in Example II, hydrogen peroxide can have an inhibitory effect on odor production by the mixed bacteria that comprise the microflora of the salivary sediment system. However, it will be difficult for peroxide to persist in this system or in dento-gingival or tongue plaque *in situ* because their microfloras contain bacteria that have exceptionally high catalase and peroxidase activities. (Ryan and Kleinberg (1995) *Archs. oral Biol.* 40:743-752). Consequently, for peroxide to be effective in the mouth, it is necessary to inhibit this catalase activity. Otherwise, peroxide levels well over 1% would be necessary to reduce the oral malodor and this concentration can be harmful to the oral soft tissues. An agent presently documented to be inhibitory of catalase activity is chloride ion.

This assay examined the effects of rinsing with a combination of $ZnCl_2$, H_2O_2 and NaCl on the E_h and odor producing activity of the oral bacteria. The oral rinse consisted of $ZnCl_2$ at 6 mM (0.08%), NaCl at 500 mM (2.9%) and H_2O_2 at 1%.

The oral rinse was tested for its effects on the E_h lowering and oral malodor forming activity of the oral bacteria. The oral malodor formation was assessed organoleptically, measuring VSC using a Halimeter (Model RH-17A Interscan Portable Analyzer) and measuring the formation of indole/skatole using Kovac's method. Gadebusch H.H. and Gabriel S. (1956). "Modified Stable Kovac's Reagent for the Detection of Indole", *Amer. J. Clin. Path.* 26, 1373-1375. The organoleptic measurements were done by having a trained individual smell the odor produced and rate the odor on a scale of 0-4, with zero indicating no malodor and four indicating strong malodor. In the Kovac method, Kovac's Reagent (P-Dimethylaminobenzaldehyde dissolved in amyl alcohol and acidified with HCl) was added to each assay sample and the bluish/red color was measured at 567 nm in a spectrophotometer.

Experiments were run on subjects who had fasted and carried out no oral hygiene for the last 12 h. Testing was started between approximately 9 a.m. and 10 a.m. For the baseline sample, the subject cleared saliva from the mouth by gentle expectoration and rested for a period of 2 minutes to allow fresh saliva to collect. This saliva was then spit into a test tube chilled in cracked ice. If the collected saliva was less than 750 ml a further 2 minute period was used for collection. Following the baseline collection, the subject rinsed his/her mouth with 5 ml of the test solution for 20 seconds and a saliva sample was collected as above. The collection once more was for a 2 minute period. Saliva samples were also

collected at different times up to four hours following the rinse. 750 ml of each saliva sample was immediately incubated in a 37°C water bath and the E_h , odor index, and volatile sulfur compounds (VSC) in each sample were determined at 0, 15 and 30 minutes and at 1, 2, 3, 4, 8 and 24 h; these values were averaged to give a mean value assay of each parameter in each sample. Subjects were tested on 3 different days. Day one was without the use of a rinse, day two with a water rinse and day three with a zinc-peroxide-chloride ion rinse.

Figure 4 shows the results after the zinc-hydrogen peroxide-chloride ion rinsing. As shown by this figure, the zinc-hydrogen peroxide-chloride ion composition of the present invention favored a more positive E_h level and its effect on odor was dramatic. Odor production determined organoleptically and by indole/skatole formation were almost totally retarded and the production of VSC was significantly reduced. No changes from base-line (BL) were seen when water was the rinsing solution or no rinsing was done. The oral composition of the present invention affected the E_h positively and suppressed odor parameters even after 4 hours following the rinse. Only indole/skatole formation began to return after 4 hours. These results demonstrate the ability of the oral compositions described herein to reduce oral malodor by preventing the ability of the oral bacteria to reduce the E_h of the oral cavity.

Example IV

The ability of a zinc chloride rinse solution to reduce or prevent the formation of volatile sulfur compounds (VSC) *in vivo* following successive challenges with a cysteine rinse was studied in this Example. The rinse solution consisted of zinc chloride at 6 mM (0.08%). Experiments were run on subjects who had fasted and carried out no oral hygiene for at least 12 hours. The subject rinsed his/her mouth with 5 ml of a 6 mM cysteine solution for 30 seconds to stimulate substantial VSC production by the oral bacteria. This test is analogous to the use of glucose as a challenge substrate to determine the glycolytic activity of dental plaque bacteria (Stephan, R.M. 1944. *J. dent. Res.* 23, 257) or ingestion of a fixed amount of glucose as a challenge in assaying for diabetes activity. Before and following the rinse, a Halimeter instrument was used to measure the volatile sulfur compounds (VSC) produced. Simultaneously, a platinum electrode was placed on the posterior dorsum surface of the tongue, just anterior to the circumvallate papillae to measure the E_h . After 20 minutes, the subject rinsed his/her mouth with 5 ml of the zinc chloride solution for 30 seconds. Again, the VSC of the subject's mouth air and the E_h were measured. The subject then rinsed successively with the 6 mM cysteine solution at 40, 60, 140, 320 and 380 minutes and the VSC and E_h of the

subject's oral cavity were re-measured each time (see Figure 5) using the Halimeter and the platinum electrode. As shown in Figure 5, the zinc chloride rinse reduced the ability of the oral bacteria to produce the VSC response to cysteine rinsing and retarded the production of the VSC response to subsequent cysteine rinses for about 5 hours thereafter. The same figure also shows the E_h response that occurs co-incident with the VSC response after cysteine challenging. The E_h was raised to a small degree during rinsing with the zinc chloride solution. Clearance of the cysteine by saliva and rapid utilization of the cysteine by the oral bacteria facilitates the removal of the cysteine and the returning of the VSC and to their respective baselines during each cysteine challenge episode. As demonstrated by this *in vivo* assay, zinc chloride is a compound that is able to reduce the ability of the oral bacteria to generate volatile sulfur compounds. This contributes to reduced oral putrefaction activity and in turn reduced oral malodor production and gingivitis-periodontitis.

Example V

The ability of a zinc-hydrogen peroxide-salt composition to reduce or prevent the formation of VSC *in vivo* following successive challenges with a cysteine rinse was studied in this Example in the same way as in Example IV. The oral rinse consisted of zinc chloride at 6 mM (0.08%), hydrogen peroxide at 1% and sodium chloride at 500 mM (2.9%). As before, 5 ml of the zinc/peroxide/chloride ion composition of this invention was applied as a rinse for 30 seconds. VSC and E_h measurements were also carried out as before and the results are shown in Figure 6. The zinc-peroxide-chloride ion rinse of this invention showed greater inhibition of VSC production and for a longer time than in Example IV where the zinc chloride was tested without peroxide and chloride ion. The E_h rose strongly and more with the zinc chloride-peroxide-sodium chloride composition following its application than did zinc chloride alone in Example IV and the subsequent E_h responses were initially inhibited and return to the baseline response was delayed. This Example shows that the oral composition of the present invention is able to significantly inhibit VSC production from cysteine and its ability to lower the E_h , and has the ability to raise the E_h when introduced *in vivo*.

Example VI

The ability of a zinc-chloride-sodium chlorite composition to reduce or prevent the formation of VSC *in vivo* following successive challenges with a cysteine rinse was studied in this Example in the same way as in Examples IV and V. The oral rinse consisted of zinc

chloride at 6 mM (0.08%) and sodium chlorite at 0.5%. As before, 5 ml of the zinc/peroxide/chloride ion composition of this invention was applied as a rinse for 30 seconds. VSC and E_h measurements were also carried out as before and the results are shown in Figure 7. The zinc-chloride-sodium chlorite rinse of this invention showed as in Example V, greater inhibition of VSC production and for a longer time than in Example IV where the zinc chloride was tested without sodium chlorite. The E_h rose strongly and more with the zinc chloride-sodium chlorite composition following its application than did zinc chloride alone in Example IV; the subsequent E_h responses were initially inhibited and return to the baseline response was delayed. This Example shows that the oral composition of the present invention is able to significantly inhibit VSC production from cysteine and its ability to lower the E_h , and has the ability to raise the E_h upon introduction into the oral cavity.

Example VII

The abilities of 0.1% sodium chlorite rinse solutions, one without and one with zinc chloride at 6 mM (0.08%) were compared in this Example for their abilities to reduce or prevent the formation of VSC *in vivo* following successive challenges with a cysteine rinse as before. The oral rinse consisted of application of 5 ml for 30 seconds of either an 0.1% sodium chlorite solution or an 0.1% sodium chlorite solution to which zinc chloride at 0.08% had been added. VSC and E_h measurements were carried out as before and the results are shown in Figures 8 and 9. The sodium chlorite showed no inhibition of VSC production (Figure 8) whereas the zinc chloride-sodium chlorite combination was inhibitory and more so than observed for zinc chloride alone (Figure 5) in Example IV. This Example shows that combining zinc chloride and sodium chlorite can produce a greater effect than expected from simple addition of their individual effects.

We claim:

- 1 1. An oral composition comprising a zinc compound containing freely
2 available zinc ions, wherein the concentration of freely available zinc ions ranges from about
3 0.02% to about 0.2% and at least one stabilized E_h raising compound, wherein the concentration
4 of the E_h raising compound ranges from about 0.1% to about 3.0%, distributed in an oral
5 vehicle, wherein the zinc compound and the E_h raising compound are stored separately and
6 mixed prior to using the oral composition.
- 1 2. The oral composition of claim 1 wherein the concentration of the freely
2 available zinc ranges from about 0.04% to about 0.12% by weight.
- 1 3. The oral composition of claim 1 wherein the concentration of the E_h
2 raising compound ranges from about 0.1% to about 1.0% by weight.
- 1 4. The oral composition of claim 1 wherein the zinc compound containing
2 freely available zinc ions is selected from the group consisting of zinc chloride, zinc sulfate,
3 zinc acetate, zinc lactate, zinc salicylate, and zinc nitrate.
- 1 5. The oral composition of claim 1 wherein the E_h raising compound is
2 selected from the group consisting of hydrogen peroxide, oxyhalogen, methylene blue and
3 sodium chloride.
- 1 6. The oral composition of claim 1 wherein the zinc compound is zinc
2 chloride and the E_h raising compound is hydrogen peroxide and the composition further
3 comprises a chloride ion compound, wherein the chloride ion concentration ranges from about
4 0.5% to about 2.5% by weight.
- 1 7. The oral composition of claim 6 wherein the pH of the zinc compound
2 during storage ranges from about 3.0 to about 6.0 and the pH of the hydrogen peroxide during
3 storage ranges from about 3.0 to about 6.0.
- 1 8. The oral composition of claim 6 wherein the chloride ion compound is
2 sodium chloride.

1 9. The oral composition of claim 1 wherein the zinc compound is zinc
2 chloride and the E_h raising compound is sodium chlorite.

1 10. The oral composition of claim 9 wherein the pH of the zinc compound
- 2 during storage ranges from about 3.0 to about 6.0 and the pH of the sodium chlorite during
3 storage ranges from about 7.0 to about 8.5.

1 11. The oral composition of claim 10 wherein the composition further
2 comprises a chloride ion compound.

1 12. The oral composition of claim 11 wherein the chloride ion compound
2 is sodium chloride.

1 13. The oral composition of claim 1 wherein the oral composition is a dental
2 care product, food product, lozenge, chewing gum or a confection.

1 14. A method for inhibiting the formation of sulfur containing anions in the
2 oral cavity and preventing a lowering of the E_h of the oral cavity by delivering into the oral
3 cavity a therapeutically effective amount of an oral composition comprising a zinc compound
4 containing freely available zinc ions having a concentration ranging from about 0.02% to about
5 0.2% and at least one stabilized E_h raising compound, wherein the concentration of the E_h
6 raising compound range from about 0.1% to about 3.0%, distributed in an oral vehicle, wherein
7 the zinc compound and the E_h raising compound are stored separately and mixed prior to using
8 the oral composition.

1 15. The method of claim 14 wherein the concentration of the freely available
2 zinc ion ranges from about .04% to about 0.12% by weight.

1 16. The method of claim 14 wherein the concentration of E_h raising
2 compound ranges from about 0.1% to about 1.0% by weight.

1 17. The method of claim 14 wherein the zinc compound containing freely
2 available zinc is selected from the group consisting of zinc chloride, zinc sulfate, zinc acetate,
3 zinc lactate, zinc salicylate, and zinc nitrate.

1 18. The method of claim 14 wherein the E_h raising compound is selected
2 from the group consisting of hydrogen peroxide, oxyhalogen, methylene blue and sodium
3 chloride.

1 19. The method of claim 14 wherein the zinc compound is zinc chloride and
2 the E_h raising compound is hydrogen peroxide and the composition further comprises a chloride
3 ion compound, wherein the chloride ion concentration ranges from about 0.5% to about 2.5%
4 by weight.

1 20. The method of claim 19 wherein the pH of the zinc compound during
2 storage ranges from about 3.0 to about 6.0 and the pH of the hydrogen peroxide during storage
3 ranges from about 3.0 to 6.0.

1 21. The method of claim 19 wherein the chloride ion compound is sodium
2 chloride.

1 22. The method of claim 14 wherein the zinc compound is zinc chloride and
2 the E_h raising compound is sodium chlorite.

1 23. The method of claim 22 wherein the pH of the zinc compound during
2 storage ranges from about 3.0 to about 6.0 and the pH of the sodium chlorite during storage
3 ranges from about 7.0 to about 8.5.

1 24. The method of claim 23 wherein the composition further comprises a
2 chloride ion compound.

1 25. The method of claim 24 wherein the chloride ion compound is sodium
2 chloride.

1 26. The method of claim 14 wherein the oral composition is a dental care
2 product, food product, lozenge, chewing gum or a confection.

1 27. A method for reducing oral malodor and gingivitis-periodontitis by
2 delivering into the oral cavity a therapeutically effective amount of an oral composition
3 comprising a zinc compound containing freely available zinc ions having a concentration
4 ranging from about 0.02% to 0.2% and at least one stabilized E_h raising compound, wherein
5 the concentration of the E_h raising compound ranges from about 0.1% to about 3.0%,
6 distributed in an oral vehicle, wherein the zinc compound and the E_h raising compound are
7 stored separately and mixed prior to using the oral composition.

1 28. A method for claim 27 wherein the concentration of the freely available
2 zinc ion ranges from about .04% to about 0.12% by weight.

1 29. A method for claim 27 wherein the concentration of E_h raising
2 compound ranges from about 0.1% to about 1.0% by weight.

1 30. A method for claim 27 wherein the zinc compound containing freely
2 available zinc is selected from the group consisting of zinc chloride, zinc sulfate, zinc acetate,
3 zinc lactate, zinc salicylate, and zinc nitrate.

1 31. A method for claim 27 wherein the E_h raising compound is selected from
2 the group consisting of hydrogen peroxide, oxyhalogen, methylene blue and sodium chloride.

1 32. A method for claim 27 wherein the zinc compound is zinc chloride and
2 the E_h raising compound is hydrogen peroxide and the composition further comprises a chloride
3 ion compound, wherein the chloride ion concentration ranges from about 0.5% to about 2.5%
4 by weight.

1 33. A method for claim 27 wherein the pH of the zinc compound during
2 storage ranges from about 3.0 to about 6.0 and the pH of the hydrogen peroxide during storage
3 ranges from about 3.0 to 6.0.

1/9




FIG. 1

1 34. A method for claim 32 wherein the chloride ion compound is sodium
2 chloride.


1 35. A method for claim 27 wherein the zinc compound is zinc chloride and
2 the E_n raising compound is sodium chlorite.

1 36. A method for claim 35 wherein the pH of the zinc compound during
2 storage ranges from about 3.0 to about 6.0 and the pH of the sodium chlorite during storage
3 ranges from about 7.0 to about 8.5.

1 37. A method for claim 36 wherein the composition further comprises a
2 chloride ion compound.

1 38. A method for claim 37 wherein the chloride ion compound is sodium
2 chloride.

1 39. A method for claim 27 wherein the oral composition is a dental care
2 product, food product, lozenge, chewing gum or a confection.



— Zn⁺ —+— H₂O₂⁺ (0.6%) —x— Zn⁺ + H₂O₂⁺ (0.5%) —o— Supernatant plus sediment (control)
* present as ZnCl₂ at 6 mM (0.08%)
** present at 0.5%

FIG. 3

2/9




FIG. 2

5/9




FIG. 5






FIG. 4

* 10 ml FOR 30 SECONDS
 ** 24 HR MEAN
 *** 24 HR PRODUCTION
 BL-Baseline



--- cys or test rinse applied
FIG. 7

6/9



--- cys or test rinse applied

FIG. 6




FIG. 9




FIG. 8

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US97/19598

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61C 5/00; A61K 31/54, 33/14, 33/30, 33/40

US CL : 424/614, 613, 642, 665, 680; 433/215; 514/224.8

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/614, 613, 642, 665, 680; 433/215; 514/224.8

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
NoneElectronic data base consulted during the international search (name of data base and, where practicable, search terms used)
CAS on-line

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y, P	Biosis Abstract, Abstract No. 97:450683, WALTER, S. M. 'The effect of some metal ions on volatile sulfur-containing compounds originating from the oral cavity', abstract, Acta Odontologica Scandinavica. 1997, Vol. 55, No. 4, pages 261-264, see entire abstract.	1-39

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

Special categories of cited documents:	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
document defining the general state of the art which is not considered of particular relevance	"X"	document of particular relevance, the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
document published on or after the international filing date	"Y"	document of particular relevance, the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"G"	document member of the same patent family
document referring to an oral disclosure, use, exhibition or other means		
document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

19 DECEMBER 1997

Date of mailing of the international search report

10 FEB 1998

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
P.O. Box 1000

Authorized officer

A. A. F. I.

This Page Blank (uspto)